

PATIENT'S LEAFLET.
READ CAREFULLY BEFORE USING THIS MEDICINE

BURNAL OINTMENT

COMPOSITION:

Euflavine (neutral acriflavine) 0.1%

THERAPEUTIC ACTIVITY:

Antimicrobial for local treatment of infected wounds, burns or ulcers. Also it can help prevent development of local infection in wounds and burns.

WHEN SHOULD THE PREPARATION NOT BE USED?

- Do not use this medicine without consulting a doctor or a pharmacist before starting treatment if you are pregnant or breastfeeding.
- Do not use this medicine if you are sensitive to any of its ingredients.

WARNING

If you are sensitive to any type of food or medicine, inform your doctor or pharmacist before commencing treatment with this medicine.

DRUG INTERACTIONS

If you are using another drug concomitantly or if you have just finished treatment with another medicine, inform your doctor or pharmacist, in order to prevent hazards or lack of efficacy arising from drug interactions especially with other skin preparations.

SIDE EFFECTS

In addition to the desired effect of the medicine, adverse reactions may occur during the course of using this medicine such as: pruritus, burning sensation, rash or irritation.

If these side effects continue and /or are bothersome consult your doctor.

Effects that require special attention:

In the event that you experience side effects not mentioned in this leaflet, or if there is a change in your general health, consult your doctor immediately.

DOSAGE

According to doctor's or pharmacist's instructions, or according to the directions for use in this leaflet.

Do not exceed the recommended dosage.

ATTENTION !

Do not swallow!

This medicine is intended for external use only.

DIRECTIONS FOR USE

Sterile measures should be taken when applying Burnal ointment and it should be spread directly onto the burn wounds after they have been thoroughly cleansed.

The ointment should be applied once or twice daily in a sufficient layer over the entire burn area according to instructions. The affected area must be always covered by ointment. A cover dressing may or may not be used. If used, it should be changed daily or on alternate days. If for any reason the ointment is rubbed off, apply a new layer. Do not open the container except when in use.

AVOID POISONING!

This medicine, and all other medicines, must be stored in a safe place out of the reach of children and /or infants to avoid poisoning. If a child has accidentally swallowed the medicine, proceed immediately to a hospital emergency room and bring the package of the medicine with you.

Do not induce vomiting unless explicitly instructed to do so by a doctor !

Do not use medicines in the dark! Check the label and the dose each time you use your medicine. Wear glasses if you need them.

STORAGE

- Store between 15-25C

- Keep out of the reach of children.

Even if kept in their original container and stored as recommended, medicines may be kept for a limited period only. Please note the expiry date of the medicine in case of doubt, consult the pharmacist who dispensed the medicine to you.

PRESENTATION:

Tube of 20 grams.



Jerusalem Pharmaceuticals Co. Ltd.
Al-Bireh – Ramallah - Palestine