

DERMASON

Cream 1%, 0.5%

Composition:

Hydrocortisone 1%, 0.5%.

Indication:

For the topical treatment of dermatological inflammation which respond to hydrocortisone.

When should the preparation not be used?

- Do not use this medicine if you are sensitive to any of its ingredients.
- Do not use this medicine for treatment of viral infections (herpes or chicken pox), fungal infections or skin tuberculosis.
- Do not use for vaginal discharges or for diaper rash.

Do not use this medicine without consulting a doctor before starting treatment:

- If you are pregnant or breast feeding.
- If you are suffering or have suffered in the past from impaired function of the immune system, diabetes (long-term).
- In children under 12 years of age.

Warnings:

- If you are sensitive to any food or medicine inform your doctor before using this medicine.
- This medicine should not be used on large areas of skin or on open sores.
- Do not allow the medicine to come into contact with the eyes. If this happen by accident, wash the eyes immediately with water.
- Special caution should be taken when using this medicine for children and teenagers. Use

of this medicine for these age groups requires doctor's follow-up.

- Stop treatment and refer to your doctor if the condition worsens, or if the symptoms continue more than 7 days, or if they go away and come back after several days. In such cases DO NOT use any medicine containing hydrocortisone, except under direction of the doctor.

Drug Interactions:

If you are taking another drug concomitantly or if you have just finished treatment with another medicine, inform the attending doctor in order to prevent hazards or lack of efficacy due to drug interactions.

Side Effects:

In addition to the desired activity of the medicine, adverse reactions may occur during the course of treatment with this medicine, such as: redness, local flushing, tingling, itching. These side effects generally disappear within a short period following adaptation to the medicine.

If these symptoms continue or are uncomfortable, see your doctor.

Side Effects Requiring Special Attention:

If you experience itching or rash or inflammation in or around the mouth (which you did not have before using this medicine) (rare): stop treatment and consult your doctor.

In the even that you experience side effects not mentioned in this leaflet, or if there is a change in your general health, consult your doctor.

Dosage & Administration:

Unless your doctor prescribes otherwise, normal dosage:

- For adults and children over the age of two is to spread on affected area 3-4 times a day for a week.
- Do not exceed the recommended dosage.
- This medicine is usually not intended for administration to children/ infants under 2 years of age – see below:

- This medicine should not be used for more than 3 weeks in children under 4 years old, especially in areas covered by the diaper.
- Unless instructed otherwise by the doctor, do not use a bandage, which seal the affected area (plastic diapers constitute a sealing bandage).
- If using the medicine in the diaper area, do not close tightly and do not use plastic pant.
- Do not cover the affected area, unless directed to do so by the doctor.
- Spread on the affected area according to doctor's instruction.

Note:

Do not swallow! This medicine is intended for external use only.

Avoid contact of the medicine with the eyes, mucous membranes (mouth, nose, ears – see warnings).

Overdose:

If a child has accidentally swallowed the medicine, proceed immediately to a hospital emergency room and bring the package of the medicine with you.

Presentation: Tubes of 15 g.

Storage:

- Store between 15 – 25 °C.
- Keep out of reach of children.
- Discard outdated medicine.



Jerusalem Pharmaceuticals Co. Ltd.

Al-Bireh - Ramallah - Palestine