


PATIENT'S LEAFLET.  
READ CAREFULLY BEFORE USING THIS MEDICINE  
THE DISPENSING OF THIS MEDICINE REQUIRES  
A PHYSICIAN'S PRESCRIPTION.



**Composition:**

Fusylin Cream Contains:  
Fusidic acid 2%  
Fusylin Ointment Contains:  
Sodium Fusidate 2%

**Classification:**

Antibiotic / Antimicrobial.

**For What Is Fusylin Used?**

Fusylin preparations are used as topical antibacterials in treating infected skin and wounds and other conditions recommended by your doctor.

**This Drug Should Not Be Used If:**

You are allergic to the preparation or to any of its ingredients.

**Inform Your Doctor Before Using This Drug If:**

- You have ever had an allergic reaction to fusidic acid or its salts, or to any other drug, food or any substance.
- You are pregnant or breast-feeding.

**Drug Interactions:**

Although there are no known interactions when fusidic acid or its salts are used topically, it is wise to inform your doctor or pharmacist if you are taking or using externally other drugs, or you have recently stopped treatment with another drug.

**Side Effects:**

Along with its needed effects, a medicine may cause unwanted effects such as rare mild irritation, pain (when medication is applied to deep leg ulcers), allergic reaction (swelling, rash and itching). If these side effects continue and become bothersome, check with your doctor.

**Proper Use Of This Medicine:**

- Check the expiry date. Do not use if expired.
- Use exactly as prescribed by your doctor and follow your pharmacist's directions.

**Precautions While Using This Medicine:**

- This drug is for external use only.
- Do not give this drug to anyone else to avoid causing any harm.
- Do not by yourself interrupt the period of treatment prescribed to you.
- Do not repeat the same medicament and do not increase doses without consulting your doctor.
- Avoid Contacts with eyes.

**Poisoning:**

Poisoning has not been known to occur during topical therapy with Fusylin. If you or a child or anyone else has accidentally swallowed Fusylin, proceed immediately to a hospital emergency room and bring the medicine with you.

**Directions For Use:**

Use according to your doctor's or pharmacist's instructions only. Fusylin ointment or cream is applied to the lesion 3 or 4 times daily for a period specified by your doctor.

Whenever the lesion is to be covered with a gauze dressing, less frequent applications may be used.

**Storage:**

- Store at 15 - 25C.
- Store out of the reach of children.

**Note:**

If you need any other information not found in this leaflet, ask your doctor or pharmacist.

**Presentation:**

- Fusylin cream: tube of 15 grams.
- Fusylin ointment: tube of 15 grams.



**Jerusalem Pharmaceuticals Co. Ltd.**  
**Al-Bireh – Ramallah – Palestine**