



Composition:

Fusylin-H Cream Contains:
Fusidic Acid 2%
Hydrocortisone Acetate 1%.

Classification:

Topical Antibiotic / Corticosteroid.

For What Is Fusylin-H Used?

Fusylin-H preparation is used as topical antibacterial in treating infected skin and wounds; in addition, Fusylin-H also has anti-inflammatory activity. And other conditions recommended by your doctor.

When This Drug Should Not Be Used:

Fusylin-H is contraindicated in patients with hypersensitivity to fusidic acid, hydrocortisone acetate, or other components of the cream. As with other topical antibiotic/corticosteroid combination preparation, Fusylin-H is contraindicated in bacterial infections due to non-susceptible organisms, fungal infections, tuberculosis of the skin, syphilitic skin infections, chicken pox, eruptions following vaccinations and viral diseases of skin in general.

Inform Your Doctor Before Using This Drug If:

- You have ever had an allergic reaction to fusidic acid, hydrocortisone, or to any other drug, food or any substance.
- You are pregnant or breast-feeding.
- In children under 3 years of age.

Drug Interactions:

Although there are no known interactions when Fusylin-H are used topically, it is wise to inform your doctor or pharmacist if you are taking or using externally other drugs, or you have recently stopped treatment with another drug.

Side Effects:

Along with its needed effects, a medicine may cause unwanted effects such as rare mild irritation, pain (when medication is applied to deep leg ulcers), allergic reaction (swelling, rash and itching). If these side effects continue and become bothersome, check with your doctor.

Proper Use Of This Medicine:

- Check the expiry date. Do not use if expired.
- Use exactly as prescribed by your doctor and follow your pharmacist's directions.

Precautions While Using This Medicine:

- This drug is for external use only.
- Do not give this drug to anyone else to avoid causing any harm.
- Do not interrupt the period of treatment prescribed to you by yourself.
- Do not repeat the same medicament and do not increase doses without consulting your doctor.

- Avoid Contacts with eyes.
- Stop using this medicine if there is no improvement within 14 days.

Poisoning:

Poisoning has not been known to occur during topical therapy with Fusylin-H. If you or a child or anyone else has accidentally swallowed Fusylin-H, proceed immediately to a hospital emergency room and bring the medicine with you.

Directions For Use:

Use according to your doctor or pharmacist's instructions only. Fusylin-H cream is applied to the lesion 3 times daily and gently massaged into the affected areas for a period specified by your doctor. Whenever the lesion is to be covered with gauze dressing, less frequent applications may be used.

Storage:

- Store at 15° – 25°C.
- Keep out of the reach of children.

Note:

If you need any other information not found in this leaflet, ask your doctor or pharmacist.

Presentation:

Fusylin-H cream: tube of 15 grams.



Jerusalem Pharmaceuticals Co. Ltd.

Al-Bireh – Ramallah - Palestine