



Composition:

Silver Sulfadiazine 1%

Indications:

Antimicrobial preparation for local treatment of burns, infected pressure sores and leg ulcers.

When Should the Preparation not be used?

- Do not use this medicine if you are sensitive to any of its ingredients.
- Do not use on premature infants during the first few months of life.

Do Not Take This Medicine Without Consulting A Doctor Before Starting Treatment:

- If you are pregnant or breastfeeding.
- If you have a deficiency of the G₆PD enzyme (Sensitivity to Fava bean).
- If you suffer or have suffered in the past from impaired function of the liver or kidney.

Warnings:

- During treatment with this medicine the following tests should be performed: blood, urine and kidney function.
- If you are sensitive to any type of food or medicine, inform your doctor before commencing treatment with this medicine.

Drug Interactions:

You have to inform your doctor if you are taking another drug at the same time or you have recently stopped treatment with another drug. This is to avoid problems that may arise from drug interactions especially with: other skin preparations containing proteolytic enzymes.

Side Effects:

In addition to the desired effect of the medicine, adverse reactions may occur during the course of taking this medicine, for example: pruritus or burning sensation.

If these side continue and/or bothersome consult your doctor.

Effects that require special attention:

Hypersensitivity to sunrays, local irritation, rash or inflammation (which were not present before use of the preparation).

When the preparation is used over extensive areas it may be absorbed into the vascular system. If this occurs adverse reactions may appear that are characteristic of an oral dose of the preparation: fever, sore throat, tiredness or extreme weakness (rare): stop treatment and consult your doctor.

In the event that you experience side effects not mentioned in this leaflet, or if there is a change in your general health, consult your doctor immediately.

Dosage:

- According to doctor's instructions only.
- Do not exceed the recommended dose.

Note:

- Do not swallow! This medicine is intended for external use only.
- Do not take medicines in the dark!
- Check the label and the dose each time you take your medicine. Wear glasses if you need them.

Directions for Use:

Sterile measure should be taken when applying Silvaderm cream and it should be spread directly onto the burn wounds after they have been thoroughly cleansed.

The cream should be applied in a layer 3-5 mm thick, over the entire burn area, according to doctor's instructions. The affected area must be always covered by cream. If for any reason the cream is rubbed off, apply a new layer. A separate tube should be used for each patient. Do not open the container except when in use.

Missed Dose:

If you miss a dose of this medicine, take it as soon as you remember. However, if it is almost time for your next dose, skip the missed dose and go back to your regular dosing schedule. Do not double doses.

Overdose:

If you have taken an overdose or if a child has accidentally swallowed the medicine, proceed immediately to a hospital emergency room and bring the package of the medicine with you.

Presentation:

- Tube of 50 g.
- Jars of 75 , 200 & 500 g.

Storage Conditions:

- Store below 25 °C.
- Keep out of reach of children.
- Protect from light.



Jerusalem Pharmaceuticals Co. Ltd.

Al-Bireh - Ramallah - Palestine