

# BETA-CANAZOLE Cream

## Composition:

Each tube of cream contains:

Clotrimazole	1%
Betamethasone	0.05%.

## Therapeutic Activity:

This product is indicated for the treatment of skin inflammations involving fungal infection.

## WHEN SHOULD THE PREPARATION NOT BE USED?

- Do not use this medicine if there known hypersensitivity to one of its ingredients.

## ***Do Not Take This Medicine Without Consulting A Doctor Before Starting Treatment:***

- If you are pregnant or breastfeeding.

## Warnings:

- Do not use occluded bandaging unless otherwise instructed by the doctor (nylon diapers are considered occluded bandaging).
- Do not allow contact with the eyes. In case of such contact, wash immediately with water.
- If you are sensitive to any material or drug, inform your doctor prior to commencing treatment with this product.
- Do not use this product for any purpose other than that it was indicated.

- Do not use this product on extensive skin areas or treatment of the face should be open wounds carried out with caution and for a short period only.
- Special caution is required when using this product in children. Use of this product in children and in adolescents should be under medical supervision.

## Drug Interactions:

You have to inform your doctor if you are taking another drug at the same time or you have recently stopped treatment with another drug. This is to avoid problems that may arise from drug interaction especially with: another drug, consult your doctor to avoid risks or inefficacy arising from drug interactions.

## Side Effects:

In addition to the desired effect of the medicine, adverse reactions may occur during the course of taking this medicine, for example: rash, irritation, itching, dryness, redness, sense of heat and tingling.

These side effects usually disappear shortly after adaptation to the product.

If you feel side effects not mentioned in this leaflet, or if there has been any change in your general health, consult your doctor immediately.

## Dosage & Administration:

- Apply a thin layer of Beta-Canazole on the affected area, twice daily.
- Dosage according to the doctor's instructions only.
- Do not exceed the recommended dose or treatment period.

## Dosage for Children:

- In case of treating an area covered with a diaper, do not tighten the diaper and do not use nylon pants.
- In infants and children under the age of 4, do not use this product for more than 3 weeks.
- If your condition has not improved after a few days, contact your doctor again.

## Note:

- Do not swallow! This product is for external use only.
- Avoid contact of this drug with the eyes and mucous membranes (i.e. mouth and nose).
- Complete the course of treatment recommended by your doctor.
- Even if your condition has improved, do not discontinue treatment with this drug without consulting your doctor.
- Do not take medicines in the dark! Check the label and the dose each time you take your medicine. Wear glasses if you need them.

## Overdose:

If a child has accidentally swallowed the medicine, proceed immediately to a hospital emergency room and bring the package of the medicine with you.

## Storage Conditions:

- Store at 15-30°C.
- Keep out of reach of children.

## Presentation:

Beta-Canazole Cream: Tube of 15 grams.



**Jerusalem Pharmaceuticals Co. Ltd.**  
**Al-Bireh – Ramallah – Palestine**