

CANAZOLE Cream

Composition:

Each tube of cream contains:
Clotrimazole 1%

Therapeutic Activity:

This preparation is intended for the treatment of skin infections caused by fungi of different strains, such as: Candida, dermatophytes and other mycoses.

WHEN SHOULD THE PREPARATION NOT BE USED?

- Do not use this medicine if you are sensitive to any of its ingredients.

Do Not Take This Medicine Without Consulting A Doctor Before Starting Treatment:

- If you are pregnant or if you are breastfeeding.

Warnings:

- Do not use this preparation frequently, or for a long period, without consulting a doctor.
- Avoid contact with the eyes. In case of contact with the eyes, wash thoroughly with water.
- If you are sensitive to any type of food or medicine, inform your doctor before commencing treatment with this preparation.

Drug Interactions:

You have to inform your doctor if you are taking another drug at the same time or you have recently stopped treatment with another drug. Inform the attending doctor in order to prevent hazards or lack of efficacy arising from drug interactions.

Side Effects:

In addition to the desired effect of the medicine, adverse reactions may occur during the course of taking this medicine, for example: irritation and itching, feeling of warmth and stinging at the treated area. These side effects usually disappear within a short time following the period of adaptation to the preparation.

Effects that require special attention:

On very rare occasions symptoms of irritation and hypersensitivity was noted that required cessation of treatment.

In the event that you experience side effects not mentioned in this leaflet, or if there is a change in your general health, consult your doctor immediately.

Dosage & Administration:

Recommended Dosage Unless Otherwise Prescribed by Your Doctor:

- Twice daily treatment is recommended (morning and evening). The usually duration of treatment, for the different myotic skin infections, lasts between 2-3 weeks.
- If there is no improvement in your condition within 4 weeks, refer to your doctor.
- Complete the treatment recommended by your doctor.
- Even if there is an improvement in your condition, do not discontinue use of this preparation without consulting your doctor.
- In order to prevent recurrence of the myotic infections (especially of the feet), continue the treatment for about two weeks after the

disappearance of all the infection's symptoms.

Directions for Use:

First, clean and dry the affected area. Canazole cream should be applied to the affected area and gently rubbed in. Generally a strip of cream is sufficient to treat an area the size of the palm of your hand.

Note:

- Do not swallow! The preparation is intended for external use only.
- Avoid contact with mucous membranes (such as the mouth and nose).
- Do not cover (e.g. with bandage) the affected area, unless explicitly instructed to do so by your doctor.
- Do not take medicines in the dark! Check the label and the dose each time you take your medicine. Wear glasses if you need them.

Avoid Poisoning:

If a child has accidentally swallowed the medicine, proceed immediately to a hospital emergency room and bring the package of the medicine with you.

Storage Conditions:

- Store below 25 °C.
- Protect from heat.
- Keep out of reach of children.

Presentation:

Canazole Cream: Tube of 20 grams.



Jerusalem Pharmaceuticals Co. Ltd.
Al-Bireh – Ramallah – Palestine